



Vegetarian diet is inversely associated with prevalence of depression in middle-older aged South Asians in the United States

Yichen Jin^a, Namratha R. Kandula^b, Alka M. Kanaya^c and Sameera A. Talegawkar^a

^aDepartment of Exercise and Nutrition Sciences, Milken Institute School of Public Health, The George Washington University, Washington, DC, USA; ²Division of General Internal Medicine, Northwestern University Feinberg School of Medicine, Chicago, IL, USA; ^cDivision of General Internal Medicine, Department of Medicine, University of California, San Francisco, CA, USA

ABSTRACT

Objective: To investigate associations between a vegetarian diet and depression among South Asians in the United States.

Design: Data from 892 South Asians (age range 40–83 y, 47% women) enrolled in the Mediators of Atherosclerosis in South Asians Living in America (MASALA) study were included. A vegetarian diet was defined as no intake of meat, poultry or fish in the previous year as reported on a validated food frequency questionnaire. Depressive symptomology was assessed using the Center for Epidemiologic Studies-Depression Scale (CES-D) and depression was classified as CES-D score ≥ 16 . Multivariable logistic regression was used and covariates included age, sex, study site, education, smoking, body mass index, acculturation, intentional exercise, alcohol and energy intake, and antidepressant medication use.

Results: Our study demonstrated 43% lower odds of depression among vegetarians ($p = 0.023$).

Conclusions: Vegetarian diet was found to be inversely associated with the prevalence of depression. Longitudinal examinations confirming these findings are needed.

ARTICLE HISTORY


Received 9 August 2018
Accepted 26 March 2019

KEYWORDS

Vegetarian; diet; depression;
South Asians

Introduction

Depression is a serious and common mood disorder affecting over 350 million people worldwide (Smith 2014). It is also a major contributor to Disability Adjusted Life Years, the overall global burden of disease, and is projected to be the leading cause of disease burden by 2030 (Smith 2014; Mathers 2008). Dietary behaviors and patterns are thought to play a role in depression. A recent review of the longitudinal associations between diet quality and depressive symptoms indicated that better diet quality examined using dietary indices and scores such as the Alternative Healthy Eating Index (AHEI), Mediterranean or Tuscan diet, among others, was associated with a lower risk

CONTACT Sameera A. Talegawkar  sameera.talegawkar@alumni.tufts.edu 

(Molendijk et al. 2018). Additionally, consumption of foods such as fish and vegetables were also related to lower depression risk.

A vegetarian diet has been considered to be healthy and reported to be associated with lower risk of some chronic diseases including diabetes, hypertension and subclinical atherosclerosis as well as lower body mass index (Fraser 2009; Acosta-Navarro et al. 2017; Baines, Powers, and Brown 2007). In our previous analyses, we showed inverse associations between a vegetarian diet and visceral fat, total and LDL cholesterol, fasting glucose, insulin resistance and odds of fatty liver and any coronary artery calcium (Jin et al. 2018). However, observational research conducted to date has demonstrated inconsistent findings on the associations of a vegetarian diet with depression with the majority of studies demonstrating higher risk of depression among vegetarians. For example, using cross-sectional data from the Avon Longitudinal Study of Parents and Children, Hibbeln et al. found that the risk of depression was 1.67 times higher for vegetarian compared to non-vegetarian men (Hibbeln et al. 2018). In a cohort of American college students, vegetarians and semi-vegetarians were more likely to report being depressed than non-vegetarians (Forestell and Nezlek 2018). Also, using data from the German Health Interview and Examination Survey and its Mental Health Supplement, the prevalence of mental disorders including depression was higher among both completely and predominantly vegetarians (Michalak, Zhang, and Jacobi 2012). In contrast to these studies, a longitudinal Spanish cohort study reported that higher adherence to a pro-vegetarian dietary pattern (operationalized to quantify the habit of consuming preferentially plant-derived foods instead of animal-derived foods but without the need to follow a strict vegetarian diet) was associated with a lower risk of depression and use of antidepressant drugs (Sánchez-Villegas et al. 2015), and using data from volunteers of Seventh Day Adventist communities, researchers found that vegetarians reported less negative emotions as assessed by Depression Anxiety Stress Scale and Profile of Mood States questionnaires (Beezhold, Johnston, and Daigle 2010).

Immigrants may be vulnerable to depressive symptoms because of additional stress from cultural barriers, lack of social support (Zisberg 2017), and social companionship such as eating together which is beneficial for psychological wellbeing (Wong, Yoo, and Stewart 2007). South Asians are individuals from countries of the Indian subcontinent including Bangladesh, Bhutan, India, Maldives, Pakistan, Nepal and Sri Lanka and are one of the fastest growing minority population in the United States (US) (Hoeffel et al. 2012). A high proportion of South Asians follow a vegetarian diet due to cultural traditions and religious reasons (Singh et al. 2014). The overall objective of this study was to examine the associations between following a vegetarian diet and depression in a cohort of South Asian immigrants in the US.

Materials and methods

Participants and data collection

We used data from the baseline exam of the Mediators of Atherosclerosis in South Asians Living in America (MASALA) study, a community-based cohort of South Asians from the San Francisco Bay and the greater Chicago areas. A total of 906 men and women, without known cardiovascular disease were recruited between 2010

and 2013 using surname-based recruitment methods; 14 participants with incomplete food frequency questionnaires or implausible energy intakes were excluded, leaving the analytic sample of 892 participants for these analyses. Detailed information on the MASALA study has been provided elsewhere (Kanaya et al. 2013). The study protocol was approved by the institutional review boards of the University of California, San Francisco and Northwestern University. All participants signed informed consent prior to undergoing study procedures.

Measures

Depression was assessed using the Center for Epidemiologic Studies – Depression Scale (CES-D), which measured depressive symptoms in the previous week with a 20-item scale such as ‘I am quick tempered’ (Radloff 1977). Participants responded ‘almost never’, ‘sometimes’, ‘often’, and ‘almost always’ on the 20-item and received 0, 1, 2, and 3 points, respectively, so the overall score ranges 0–60. Participants with a score 16 or above were considered to have depression (Roux et al. 2006).

Dietary data were assessed using the validated Study of Health Assessment and Risk in Ethnic Groups food frequency questionnaire (FFQ) (Kelemen et al. 2003). A participant was classified as being vegetarian, if they reported no consumption of meat, poultry, or fish in the previous year on the FFQ.

Covariates were selected based on previous literature and univariate analysis, and included age, sex, study site, education, income, smoking status, body mass index (BMI), acculturation, intentional exercise, alcohol and energy intakes, and antidepressant medication use. Acculturation was captured using the Traditional Cultural Beliefs scale which assessed how strongly participants believed that South Asian cultural practices should be maintained in the US and was categorized into strong, moderate and weak based on established cut points (Kanaya et al. 2014). We used the Typical Week’s Physical Activity Questionnaire to assess intentional exercise (walking for exercise, dance, conditional activities, and sports), and the total metabolic equivalent minutes/week were used. Alcohol and energy intakes were estimated from the FFQ and used as grams of ethanol/day and kcal/day for analysis. The detailed measurements of covariates is available elsewhere (Kanaya et al. 2013).

Statistical analysis

Socio-demographic characteristics were compared between vegetarians and non-vegetarians with t-test or Mann–Whitney U test and Chi-square test for continuous and categorical variables, and mean (standard deviation), median (interquartile range) or percentages were reported, respectively. Multivariable logistic regression was used to examine the associations between depression and vegetarian status adjusting for covariates. In order to account for overall diet quality, a sensitivity analysis was conducted by additional adjusting for the AHEI score (Chiuve et al. 2012). The AHEI includes 11 food components with a total score ranged from 0 to 110, with higher AHEI scores indicating better diet quality. SAS 9.4 was used for all analyses with $p < 0.05$ considered to be statistically significant.

Results

Table 1 shows socio-demographic characteristics by vegetarian status. About 38% of South Asians were vegetarian. Women, non-smokers, those with strong traditional South Asian beliefs and lower BMI tended to consume a vegetarian diet ($p < 0.05$ for all). About thirteen percent of South Asians in this cohort had depression, and the prevalence of depression was higher among non-vegetarians (9.9% for vegetarians vs. 14.4% for non-vegetarians, $p = 0.049$).

In the adjusted model, vegetarians had 43% lower odds of depression than non-vegetarians (odds ratio [OR] 0.57, 95% confidence interval [CI]: 0.35–0.92, $p = 0.023$) (**Table 2**). Among the covariates, education, income and smoking status were also independently associated with depression, wherein a bachelor's degree or higher (OR = 0.53,

Table 1. Socio-demographic and health characteristics by vegetarian status in the MASALA cohort, 2010–2013.

	Total N = 892	Vegetarian N = 335	Non-vegetarian N = 557	P value
Age (y)	55.3 (9.4)	55.6 (9.2)	55.2 (9.5)	0.558
Women, %	47.1	57.6	40.8	<0.001
Current smoker, %	3.1	1.5	4.1	0.029
Education \geq Bachelor's Degree, %	87.8	88.1	87.6	0.843
Annual income \geq \$75 K, % (n = 866)	73.6	70.8	75.2	0.158
Traditional cultural beliefs, %				<0.001
- Strong	33.4	44.7	26.4	
- Intermediate	29.9	29.0	30.4	
- Weak	36.7	26.1	43.2	
Intentional exercise (MET min/week)	945 (315–1856)	945 (420–1755)	960 (315–1890)	0.99
Body Mass Index (kg/m ²)	26.0 (4.1)	25.6 (4.1)	26.2 (4.1)	0.027
Alcohol intake (grams of ethanol/day)	2.8 (5.9)	1.0 (4.3)	3.8 (6.5)	<0.001
Energy intake (kcal/day)	1682 (504)	1597 (459)	1734 (523)	<0.001
Depression ^a , %	12.7	9.9	14.4	0.049
Antidepressant use, %	3.7	3.3	4.0	0.610

Notes: Data are mean (SD), median (interquartile range) or percentage.

^aDepression defined as Center for Epidemiologic Studies – Depression Scale (CES-D) score ≥ 16 .

Table 2. Odds ratio (OR) for cross-sectional associations between vegetarian status and depression in MASALA cohort, 2010–2013.

	OR (95% CI)	P value
Vegetarian vs. non-vegetarian	0.57 (0.35–0.92)	0.023
<i>Covariates</i>		
Age, y	1.00 (0.98–1.02)	0.940
Women vs. men	1.49 (0.93–2.37)	0.095
Study site, NWU vs. UCSF	1.09 (0.70–1.69)	0.712
Bachelor's Degree and above vs. below	0.53 (0.30–0.93)	0.028
Annual income \geq \$75 K vs. <\$75K	0.46 (0.28–0.75)	0.002
Current smoker vs. never/former smoker	3.25 (1.24–8.50)	0.016
Body mass index, kg/m ²	0.98 (0.93–1.03)	0.434
Traditional cultural beliefs		
- Intermediate vs. strong	0.81 (0.48–1.38)	0.443
- Weak vs. strong	0.70 (0.40–1.23)	0.211
Intentional exercise, MET min/week	1.00 (1.00–1.00)	0.135
Alcohol intake, grams of ethanol/day	0.98 (0.94–1.02)	0.332
Energy intake, kcal/day	1.00 (1.00–1.00)	0.613
Antidepressant use vs. no use	5.31 (2.39–11.78)	<.0001

Notes: NWU = Northwestern University; UCSF = University of California San Francisco; MET = Metabolic equivalent.

95% CI: 0.30–0.93, $p = 0.028$) and family income greater than or equal to \$75,000 (OR = 0.46, 95% CI: 0.28–0.75, $p = 0.002$) were associated with lower odds of depression, while being a current smoker (OR = 3.25, 95% CI: 1.24–8.50, $p = 0.016$) was associated with higher odds of depression. The significant results remained after additional adjusting for AHEI score, and the odds of depression was reduced by 4% per 1 point increase of AHEI score (OR = 0.96, 95% CI: 0.92–0.99, $p = 0.009$, data not shown).

Discussion

In our investigation of the relationship between following a vegetarian diet and depression, we found that the prevalence of depression in the MASALA cohort at baseline was 13%. And that following a vegetarian diet was associated with significantly lower odds of depression, independent of overall diet quality. Higher educational attainment and family income were also associated with lower odds of depression.

Compared to other ethnic groups, the prevalence of depression among South Asians (men: 11%, women: 15%) was higher compared to other US race/ethnic groups. In the Multi-Ethnic Study of Atherosclerosis, the prevalence of reported depression (CES-D ≥ 16) were 7%, 8%, 13%, and 5% among non-Hispanic white, African American, Hispanic, and Chinese men, and were 12%, 15%, 28%, and 11% among non-Hispanic white, African American, Hispanic, and Chinese women, respectively (Peplinski, McClelland, and Szklo 2018). Reasons for the relatively higher prevalence of depression among South Asians could be manifold including the presence of stress due to adapting to a new culture and environment (Zisberg 2017) and as our study showed, maintenance of traditional cultural beliefs.

Previous literature has shown the protective associations of a pro-vegetarian dietary pattern with depression in a Spanish cohort with majority of middle-aged men and women over a median follow-up of 8.5 years (Sánchez-Villegas et al. 2015). While other studies have reported that a vegetarian diet is associated with higher risk of depression (Hibbeln et al. 2018; Michalak, Zhang, and Jacobi 2012) due to a lower intake of polyunsaturated fatty acids such as omega-3 fatty acids which are critical for brain functioning and decrease proinflammatory cytokines production during depression (Grosso et al. 2014). However, a vegetarian diet is generally characterized by higher intakes of grains, vegetables, nuts, beans and legumes, and therefore can be rich in antioxidant nutrients, folate, phytochemicals and fiber (Gangwisch et al. 2015).

For example, using data from the Women's Health Initiative Observational Study, higher intakes of dietary fiber were associated with lower incident depression after 3 years of follow-up (Gangwisch et al. 2015). Foods with high glycemic index tend to be low in fiber and may lead to hyperglycemia causing secretion of counter-regulatory hormones leading to anxiety and mood change (Gangwisch et al. 2015). Nutrients such as folate have also been shown to be correlated with brain function. A meta-analysis of studies examining the role of serum and dietary folate found that individuals with depression had significant less folate intake, and folate metabolism produces S-adenosyl-methionine affecting neurotransmitters that are associated with depression (Bender, Hagan, and Kingston 2017). In previous analysis of the MASALA cohort, while omega-3 fatty acid intake was lower among vegetarians, no difference was observed in the consumption of polyunsaturated fatty acids between vegetarians and non-vegetarians, and

vegetarians reported higher intakes of fiber, folate and antioxidants such as vitamin C, which may partially explain the protective role of vegetarian diet on depression in the cohort at baseline (Jin et al. 2018).

Another reason for the inverse association between a vegetarian diet and depression in this cohort could be that following a vegetarian diet is a norm related to South Asian traditions, culture and religious affiliation which may result in no bias towards this practice. In the MASALA cohort, 38% of participants were vegetarians, while in the US, only 2.4% of the population were vegetarians according to the National Health and Nutrition Examination Survey 2003–2006 (Jaacks et al. 2016). Vegetarianism in the western society may be a target of bias because it challenges the social norm of western culture (MacInnis and Hodson 2017), and this may be a contributor to the presence of depression among vegetarians in the western countries shown in the previous studies. Our analysis showed that almost half of the vegetarians had strong traditional beliefs. Since religion may be an important reason for following a vegetarian diet, this may lead to a better social support among those with similar religion, resulting in protection from depression (Park and Roh 2013).

The strengths of our study include the large community-based sample of South Asians and dietary assessment which incorporated cultural factors with South Asian specific dishes that has been previously validated among South Asians in Canada. The limitations include the cross-sectional design of the study, so no causal relationships can be concluded. There may be a likelihood for social bias for the assessment of depressive symptoms, since CES-D has not been validated in a South Asian population. And while the MASALA cohort is generally representative of South Asian immigrants in the US, its participants are middle-aged and older with high education levels, and therefore our findings may not generalize to populations with younger South Asians or those with lower educational levels.

In conclusion, vegetarian diet is considered to be a healthy dietary pattern and has been shown to be associated with a lower risk of chronic diseases. Mental health is an important and growing public health concern. Our cross-sectional analyses showed that a vegetarian diet was associated with lower odds of depression among South Asians living in the US. Further investigations on longitudinal associations and mechanistic relationships of a vegetarian diet on depression are needed.

Acknowledgements

We thank Luis A. Rodriguez from the Department of Epidemiology & Biostatistics, University of California, San Francisco for providing the Alternative Health Eating Index variables for the sensitivity analysis conducted in this study.

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

The MASALA study was supported by the National Institutes of Health [grant number R01-HL-093009]. Data collection at UCSF was also supported by NIH/NCRR UCSF-CTSI [grant numbers UL1 RR024131, UL1 TR000004].

References

- Acosta-Navarro, Julio, Luiza Antoniazzi, Adriana Midori Oki, Maria Carlos Bonfim, Valeria Hong, Pedro Acosta-Cardenas, Celia Strunz, Eleonora Brunoro, Marcio Hiroshi Miname, and Wilson Salgado Filho. 2017. "Reduced Subclinical Carotid Vascular Disease and Arterial Stiffness in Vegetarian Men: The CARVOS Study." *International Journal of Cardiology* 230: 562–566.
- Baines, Surinder, Jennifer Powers, and Wendy J. Brown. 2007. "How Does the Health and Well-being of Young Australian Vegetarian and Semi-vegetarian Women Compare with Non-vegetarians?" *Public Health Nutrition* 10 (5): 436–442.
- Beezhold, Bonnie L., Carol S. Johnston, and Deanna R. Daigle. 2010. "Vegetarian Diets are Associated with Healthy Mood States: A Cross-sectional Study in Seventh Day Adventist Adults." *Nutrition Journal* 9 (1): 26.
- Bender, Ansley, Kelsey E. Hagan, and Neal Kingston. 2017. "The Association of Folate and Depression: A Meta-analysis." *Journal of Psychiatric Research* 95: 9–18.
- Chiuev, S. E., T. T. Fung, E. B. Rimm, F. B. Hu, M. L. McCullough, M. Wang, M. J. Stampfer, and W. C. Willett. 2012. "Alternative Dietary Indices Both Strongly Predict Risk of Chronic Disease." *The Journal of Nutrition* 142 (6): 1009–1018. doi:10.3945/jn.111.157222.
- Forestell, Catherine A., and John B. Nezek. 2018. "Vegetarianism, Depression, and the Five Factor Model of Personality." *Ecology of Food and Nutrition* 57 (3): 246–259.
- Fraser, Gary E. 2009. "Vegetarian Diets: What Do We Know of their Effects on Common Chronic Diseases?" *The American Journal of Clinical Nutrition* 89 (5): 1607S–1612S.
- Gangwisch, James E., Lauren Hale, Lorena Garcia, Dolores Malaspina, Mark G. Opler, Martha E. Payne, Rebecca C. Rossom, and Dorothy Lane. 2015. "High Glycemic Index Diet as a Risk Factor for Depression: Analyses from the Women's Health Initiative." *The American Journal of Clinical Nutrition* 102 (2): 454–463.
- Grosso, Giuseppe, Fabio Galvano, Stefano Marventano, Michele Malaguarnera, Claudio Bucolo, Filippo Drago, and Filippo Caraci. 2014. "Omega-3 Fatty Acids and Depression: Scientific Evidence and Biological Mechanisms." *Oxidative Medicine and Cellular Longevity* 2014.
- Hibbeln, Joseph R., Kate Northstone, Jonathan Evans, and Jean Golding. 2018. "Vegetarian Diets and Depressive Symptoms among Men." *Journal of Affective Disorders* 225: 13–17.
- Hoeffel, Elizabeth M., Sonya Rastogi, Myoung Ouk Kim, and Shahid Hasan. 2012. *The Asian Population: 2010*. Suitland, Maryland, United States: US Department of Commerce, Economics and Statistics Administration, US Census Bureau.
- Jaacks, Lindsay M., Deksha Kapoor, Kalpana Singh, K. M. Venkat Narayan, Mohammed K. Ali, M. Masood Kadir, Viswanathan Mohan, Nikhil Tandon, and Dorairaj Prabhakaran. 2016. "Vegetarianism and Cardiometabolic Disease Risk Factors: Differences Between South Asian and US Adults." *Nutrition* 32 (9): 975–984.
- Jin, Yichen, Alka M. Kanaya, Namratha R. Kandula, Luis A. Rodriguez, and Sameera A. Talegawkar. 2018. "Vegetarian Diets are Associated with Selected Cardiometabolic Risk Factors among Middle-older Aged South Asians in the United States." *The Journal of Nutrition* 148 (12): 1954–1960.
- Kanaya, A. M., S. K. Ewing, E. Vittinghoff, D. Herrington, C. Tegeler, C. Mills, and N. R. Kandula. 2014. "Acculturation and Subclinical Atherosclerosis among US South Asians: Findings from the MASALA Study." *Journal of Clinical and Experimental Research in Cardiology* 1 (1): 102.
- Kanaya, Alka M., Namratha Kandula, David Herrington, Matthew J. Budoff, Stephen Hulley, Eric Vittinghoff, and Kiang Liu. 2013. "Mediators of Atherosclerosis in South Asians Living in America (MASALA) Study: Objectives, Methods, and Cohort Description." *Clinical Cardiology* 36 (12): 713–720.
- Kelemen, Linda E., Sonia S. Anand, Vladimir Vuksan, Qilong Yi, Koon K. Teo, Sudarshan Devanesen, and Salim Yusuf. 2003. "Development and Evaluation of Cultural Food Frequency Questionnaires for South Asians, Chinese, and Europeans in North America." *J Am Diet Assoc* 103 (9): 1178–1184.

- MacInnis, Cara C., and Gordon Hodson. 2017. "It Ain't Easy Eating Greens: Evidence of Bias Toward Vegetarians and Vegans from Both Source and Target." *Group Processes & Intergroup Relations* 20 (6): 721–744.
- Mathers, Colin. 2008. *The Global Burden of Disease: 2004 Update*. Geneva, Switzerland: World Health Organization.
- Michalak, Johannes, Xiao Chi Zhang, and Frank Jacobi. 2012. "Vegetarian Diet and Mental Disorders: Results from a Representative Community Survey." *International Journal of Behavioral Nutrition and Physical Activity* 9 (1): 67.
- Molendijk, Marc, Patricio Molero, Felipe Ortuño Sánchez-Pedreño, Willem Van der Does, and Miguel Angel Martínez-González. 2018. "Diet Quality and Depression Risk: A Systematic Review and Dose-response Meta-analysis of Prospective Studies." *Journal of Affective Disorders* 226: 346–354.
- Park, Jisung, and Soonhee Roh. 2013. "Daily Spiritual Experiences, Social Support, and Depression among Elderly Korean Immigrants." *Aging & Mental Health* 17 (1): 102–108.
- Pepplinski, Brandon, Robyn McClelland, and Moyses Szklo. 2018. "Associations Between Socioeconomic Status Markers and Depressive Symptoms by Race and Gender: Results from the Multi-ethnic Study of Atherosclerosis (MESA)." *Annals of Epidemiology* 28 (8): 535–542.
- Radloff, Lenore Sawyer. 1977. "The CES-D Scale: A Self-report Depression Scale for Research in the General Population." *Applied Psychological Measurement* 1 (3): 385–401.
- Roux, Ana V. Diez, Nalini Ranjit, Lynda Powell, Sharon Jackson, Tené T. Lewis, Steven Shea, and Colin Wu. 2006. "Psychosocial Factors and Coronary Calcium in Adults Without Clinical Cardiovascular Disease." *Annals of Internal Medicine* 144 (11): 822–831.
- Sánchez-Villegas, Almudena, Patricia Henríquez-Sánchez, Miguel Ruiz-Canela, Francisca Lahortiga, Patricio Molero, Estefanía Toledo, and Miguel A Martínez-González. 2015. "A Longitudinal Analysis of Diet Quality Scores and the Risk of Incident Depression in the SUN Project." *BMC Medicine* 13 (1): 197.
- Singh, Pramil N., Kristen N. Arthur, Michael J. Orlich, Wesley James, Anil Purty, Jayakaran S. Job, Sujatha Rajaram, and Joan Sabaté. 2014. "Global Epidemiology of Obesity, Vegetarian Dietary Patterns, and Noncommunicable Disease in Asian Indians." *The American Journal of Clinical Nutrition* 100 (Suppl_1): 359S–364S.
- Smith, Kerri. 2014. "Mental Health: A World of Depression." *Nature* 515 (7526): 181.
- Wong, Sabrina T., Grace J. Yoo, and Anita L. Stewart. 2007. "An Empirical Evaluation of Social Support and Psychological Well-being in Older Chinese and Korean Immigrants." *Ethnicity and Health* 12 (1): 43–67.
- Zisberg, Anna. 2017. "Anxiety and Depression in Older Patients: The Role of Culture and Acculturation." *International Journal for Equity in Health* 16 (1): 177.